

Specification

Medium for isolation of enteropathogenic species, especially *Shigella* and *Salmonella* in food and animal feeding stuffs, according to ISO Standards.

Presentation

	Packaging Details	Shelf Life	Storage
20 Prepared Plates 90 mm with: 21 ± 1 ml	1 box with 2 packs of 10 plates/pack. Single cellophane.	3 months	2-14 °C

Composition

Composition (g/l):	
Xylose.....	3.750
L-Lysine HCl.....	5.000
Lactose.....	7.500
Sucrose.....	7.500
Sodium chloride.....	5.000
Yeast extract.....	3.000
Phenol red.....	0.080
Sodium Deoxycholate.....	1.000
Sodium thiosulfate.....	6.800
Ammonium ferric citrate.....	0.800
Agar.....	15.000

Description /Technique

Xylose Lysine Deoxycholate Agar is a selective differential medium, suitable for the detection of pathogenic enterobacteria in food, especially *Shigella*. A modification in the original formulation of Taylor allows the medium to perform to the specifications of the ISO standards. Gram positive microbiota are inhibited by the low amount of deoxycholate, whilst *Shigella* grows. Xylose, lactose or sucrose fermentation produce acidification of the medium which is shown by the indicator surrounding the colonies turning yellow. This colour disappears after 24 hours, so readings must be carried out between 18 and 24 hours.

Sulfide production from thiosulfate is easily detected because colonies become darker, due to the ferric sulfide precipitate. Lysine decarboxylation to cadaverine may also be observed in the medium, since it produces alkalization and consequently the indicator turns red.

All these reactions allow a good differentiation of *Shigella*, which other than *Edwardsiella* and *Proteus inconstans* are the only enterobacteria that do not ferment xylose and therefore show a negative fermentation reaction. *Salmonella* does ferment xylose, but it is consumed quickly and the medium becomes alkaline due to lysine decarboxylation, which may hide the reaction. The difference between *Shigella* and *Salmonella* is that the latter colonies become darker due to ferrous sulfide precipitates, which is also a common characteristic of *Edwardsiella*. Other types of enterobacteria do not suffer this phenomenon, since acid accumulation due to lactose and sucrose fermentation is so great that it avoids pH reversion by decarboxylation and even ferrous sulfide precipitate in the first 24 hours. In the quality control appear the typical colonial aspects of Enterobacteriaceae after 24 ± 3 h of incubation at 37 ° C.

Quality control

Physical/Chemical control

Color : Red pH: 7.4 ± 0.2 at 25°C

Microbiological control

Spiral Spreading: Practical range 100 ± 20 CFU. min. 50 CFU (productivity) / 10⁴-10⁶ CFU (selectivity).

Microbiological control according to ISO 11133:2014/A1:2018.

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Aerobiosis. Incubation at 37 ± 1 °C, reading after 24 ± 3 h

Microorganism

Enterococcus faecalis ATCC® 29212, WDCM 00087

Salmonella typhimurium ATCC® 14028, WDCM 00031

Salmonella enterica ATCC® 13076, WDCM 00030

Escherichia coli ATCC® 25922, WDCM 00013

Growth

Inhibited

Good (≥ 50 %) - Cult. medium & red colonies, black center

Good (≥ 50 %) - Cult. medium & red colonies, black center

Partially Inhibited (≤ 30%). Yellow colonies

Sterility Control

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Bibliography

- ATLAS, R.M., L.C. PARK (1993) Handbook of Microbiological Media for the examination of Food. CRC Press Inc. Boca Raton.
- DOWNES, F.P. & K. ITO (2001) Compendium of Methods for the Microbiological Examination of Foods. 4th ed. APHA. Washington. DC. USA.
- HORWITZ, W. (2000) Official Methods of Analysis of the AOAC International. 17th ed. Gaithersburg. MD. USA.
- ICMSF (1978) Microorganisms in Foods 1. University of Toronto Press.
- ISO Standard 6579-1 (2017) Microbiology of food chain - Horizontal method for the detection, enumeration and serotyping of *Salmonella* - Part 1 : Detection of *Salmonella* spp.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 6340:1995 STANDARD. Water Quality - Detection of *Salmonella* spp.
- ISO 19250 Standard (2010) Microbiology of food and animal feeding stuffs.- Horizontal method for the detection of *Shigella* spp.
- PASCUAL ANDERSON, M^ªR. (1992) Microbiología Alimentaria. Díaz de Santos, S.A. Madrid.
- TAYLOR, W.J. (1965) Isolation of *Shigella*. I. Xylose Lysine Agars: New media for isolation of enteric pathogens. Am. J. Clin. Path 44:471-475.
- US FDA (Food and Drug Administrations) (1998) Bacteriological Analytical Manual 8th ed. AOAC International. Gaithersburg. MD. USA.

Storage

Storage conditions: 2-14°C

Avoid direct contact with surfaces that can freeze product.